



# Sasso Performance Objectives

Age days	Average Live Weight	Average Cumulative Feed Consumption*	Feed Conversion	Males Live Weight
14	<b>0.49</b>	0.6419	1.31	<b>0.49</b>
21	<b>0.93</b>	1.6182	1.74	<b>0.95</b>
28	<b>1.56</b>	3.1356	2.01	<b>1.62</b>
35	<b>2.33</b>	5.126	2.2	<b>2.49</b>
42	<b>3.19</b>	7.337	2.3	<b>3.45</b>
49	<b>4.04</b>	9.5344	2.36	<b>4.4</b>
56	<b>4.87</b>	11.6393	2.39	<b>5.36</b>
63	<b>5.66</b>	13.7538	2.43	<b>6.22</b>
70	<b>6.36</b>	15.8364	2.49	<b>7.06</b>
77	<b>6.99</b>	18.174	2.6	<b>7.83</b>
84	<b>7.6</b>	20.9	2.75	<b>8.59</b>

<b>91</b>	<b>8.17</b>	<b>24.2649</b>	<b>2.97</b>	<b>9.32</b>
-----------	-------------	----------------	-------------	-------------

\* Weights provided are given in lbs

Performance objectives should be viewed as goals that can be achieved with good management and environmental control.

In no way does this data constitute a warranty or guarantee of the same performance, as conditions of nutrition, management, health, flock density or physical and biological environments (weather and relief) vary from producer to producer.

<b>Females</b> Live Weight	Carcass Yield Yield %
<b>0.47</b>	Carcass yields are indicatives only as they vary a lot with nutrition and management
<b>0.9</b>	
<b>1.49</b>	
<b>2.17</b>	
<b>2.93</b>	
<b>3.68</b>	67.50%
<b>4.38</b>	68.70%
<b>5.09</b>	69.70%
<b>5.66</b>	70.50%
<b>6.15</b>	71.10%
<b>6.61</b>	71.50%

